


ISSN: 2161-6728

Volume 9, Number 3, July 2019



Advances in Molecular Imaging



ISSN: 2161-6728



www.scirp.org/journal/ami

Journal Editorial Board

ISSN 2161-6728 (Print) ISSN 2161-6752 (Online)

<http://www.scirp.org/journal/ami>

Editor-in-Chief

Prof. Orhan Nalcioglu

University of California-Irvine, USA

Editorial Board

Prof. Gjumrakch Aliev

University of Atlanta, USA

Dr. Ying Bai

Heart Flow Inc., USA

Prof. Baowei Fei

Emory University, USA

Dr. Zhong-Ping Feng

University of Toronto, Canada

Prof. Richard Hans Gomer

Texas A&M University, USA

Prof. Gultekin Gulsen

University of California-Irvine, USA

Prof. Mohammad Mojammel Al Hakim

University of Southampton, UK

Dr. Maria Kempe

Lund University, Sweden

Dr. David Sigmund Liebeskind

University of California, USA

Dr. Stefan Lorkowski

Friedrich Schiller University, Germany

Prof. Kenneth Maiese

UMDNJ-New Jersey Medical School, USA

Prof. Adalberto Merighi

University of Torino, Italy

Prof. Jean-Pierre Raufman

University of Maryland, USA

Prof. Gianfranco Risuleo

Sapienza University of Rome, Italy

Prof. Steven Alan Rosenzweig

Medical University of South Carolina, USA

Prof. Phillip Ruiz

University of Miami School of Medicine, USA

Dr. Joy Sinha

ANDalyze, Inc., USA

Prof. Maurizio Sorice

Sapienza University of Rome, Italy

Prof. Lun-Quan Sun

Central South University, China

Prof. Bin Tean Teh

National Cancer Center, USA

Prof. Masakazu Toi

Kyoto University, Japan

Prof. Horst Christian Weber

Boston University, USA

Dr. Jia Lin Yang

University of New South Wales, Australia

Table of Contents

Volume 9 Number 3

July 2019

Interest of Systematic Chest Radiography during Periodic Medical Visits in Workers

M. N. Guena, R. B. Tchouezoue, F. Z. Alapha, J. M. Amvene, J. G. Fotsing.....43

Advances in Molecular Imaging (AMI)

Journal Information

SUBSCRIPTIONS

The *Advances in Molecular Imaging* (Online at Scientific Research Publishing, www.SciRP.org) is published quarterly by Scientific Research Publishing, Inc., USA.

Subscription rates:

Print: \$59 per issue.

To subscribe, please contact Journals Subscriptions Department, E-mail: sub@scirp.org

SERVICES

Advertisements

Advertisement Sales Department, E-mail: service@scirp.org

Reprints (minimum quantity 100 copies)

Reprints Co-ordinator, Scientific Research Publishing, Inc., USA.

E-mail: sub@scirp.org

COPYRIGHT

Copyright and reuse rights for the front matter of the journal:

Copyright © 2019 by Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>

Copyright for individual papers of the journal:

Copyright © 2019 by author(s) and Scientific Research Publishing Inc.

Reuse rights for individual papers:

Note: At SCIRP authors can choose between CC BY and CC BY-NC. Please consult each paper for its reuse rights.

Disclaimer of liability

Statements and opinions expressed in the articles and communications are those of the individual contributors and not the statements and opinion of Scientific Research Publishing, Inc. We assume no responsibility or liability for any damage or injury to persons or property arising out of the use of any materials, instructions, methods or ideas contained herein. We expressly disclaim any implied warranties of merchantability or fitness for a particular purpose. If expert assistance is required, the services of a competent professional person should be sought.

PRODUCTION INFORMATION

For manuscripts that have been accepted for publication, please contact:

E-mail: ami@scirp.org

Interest of Systematic Chest Radiography during Periodic Medical Visits in Workers

Mathurin Neossi Guena^{1,2*}, Romai Batchago Tchouezoue¹, Florent Zilbinkai Alpha², Jérémie Mbo Amvene³, Joseph Gonsu Fotsing^{4,5}

¹Department of Biomedical Sciences, Faculty of Sciences, University of Ngaoundéré, Ngaoundéré, Cameroon

²Imaging service, Ngaoundéré Regional Hospital, Ngaoundéré, Cameroon

³Faculty of Medicine and Biomedical Sciences, University of Ngaoundere, Garoua, Cameroon

⁴Faculty of Medicine and Biomedical Sciences, University of Yaounde I, Yaounde, Cameroon

⁵Yaounde Gynaecology Obstetrics and Pediatrics Hospital, Yaounde, Cameroon

Email: *mneossiguena@yahoo.fr

How to cite this paper: Guena, M.N., Tchouezoue, R.B., Alpha, F.Z., Amvene, J.M. and Fotsing, J.G. (2019) Interest of Systematic Chest Radiography during Periodic Medical Visits in Workers. *Advances in Molecular Imaging*, 9, 43-52.

<https://doi.org/10.4236/ami.2019.93004>

Received: April 28, 2019

Accepted: June 14, 2019

Published: June 17, 2019

Copyright © 2019 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Background: Systematic chest X-ray is the most prescribed examination by occupational physicians during periodic medical check-ups in our context, unlike in most countries where this practice has been discontinued. **Objective:** The objective was to determine the diagnostic yield and cost-effectiveness of chest radiography during periodic medical check-ups of workers in Ngaoundere. **Materials and Methods:** A cross-sectional and descriptive study was carried out in Ngaoundere Regional Hospital from January to December 2018, concerning all persons coming to perform a systematic chest X-ray in the context of the periodic medical check-up having been consented to participate in the study. Posteroanterior (PA) chest radiographs were obtained from all workers during this periodic medical check-up. The variables studied were: age, sex, clinical manifestations, antecedents, job type, cost, and chest X-ray results. Statistical analysis was performed using the Sphinx Plus²-Lexica-V5 Edition software. **Results:** 753 workers were selected for this study, of whom 88.04% were men and 11.96% were women, a sex ratio H/F is of 7.4, the average age was 39 ± 7.89 years with extremes of 24 and 56 years, the most frequent job type was the administration (38.37%), the storekeepers (17.92) and the drivers (15.27%), most of them were asymptomatic (97.47%), some had cough (2.52%), chest pain (2.12%), dyspnea (0.26%) and fever (0.13%). History was dominated by lung infection (2.39%), pleurisy (1.06%) and tuberculosis (0.79%). 7.43% of workers occasionally smoked tobacco and consumed alcohol. 41 cases of pathologies (5.44%) were found on the chest X-ray, including 1 case (0.13%) of evolutionary pathology, 40 cases (5.31%) of sequelae pathologies. When the results of the chest X-ray are compared with the clini-

cal signs and the pathological history of the workers, it is found that the only case with evolutionary abnormalities on the chest X-ray showed clinical signs and that the other cases with sequelae abnormalities had either clinical signs or antecedents related to these abnormalities. The cost of a chest X-ray at the time of our study was 5,000 FCFA (\$8.59), for a total of 3,765,000 FCFA (\$6473.74) for all the X-rays performed. **Conclusion:** In view of the low rate of abnormalities on the chest X-ray and the sequelae nature of the pathologies, it can be said that the routine X-ray of the chest during the periodic medical check-ups has a low diagnostic yield and a low cost-effectiveness ratio and is not profitable. It should be non-systematic and case-by-case based on the clinical context and background of workers.

Keywords

Chest X-Ray, Workers, Periodic Medical Check-Ups, Diagnostic Value, Cost

1. Introduction

The periodic medical examination is a visit to which the employee is subjected in the course of his employment, carried out by the working doctor, on which the company depends. This is the central element of worker health surveillance. Prescribed by law, its purpose is to pronounce on the suitability or incapacity of the worker at the job type and to ensure that he does not present any contagious disease that could endanger the health of other workers [1]. During these periodic medical examinations, the occupational doctor carries out an interrogation, the clinical examination, and can carry out or prescribe the additional examinations necessary for the determination of the compatibility between the job type and the state of health of the worker, including the detection of conditions that may lead to a contre-indication to this job, an occupational disease that may result from the professional activity of the worker and the detection of dangerous diseases for the professional environment of the worker. The regulation does not specify the examinations to be performed during the periodic medical visits, thus leaving the choice to the doctor to prescribe the examinations according to the risk and the epidemiological context [2]. Of all the complementary examinations, the chest radiography is certainly the most prescribed by the occupational physicians. Its usual justification outside any occupational exposure is the search for undifferentiated tuberculosis [3]. Adekol *et al.* in a study on routine chest X-ray in pre-employment medical examination for healthcare workers found a single abnormality that was cardiomegaly in 4.8%, so 66% was over 41 years old therefore significantly associated with age, he therefore concluded that this practice was not necessary and should be reserved for older workers [4]. Moifo *et al.* in a study assessing the role of routine chest X-ray in asymptomatic students during university enrollment in a tuberculosis-endemic area find that this practice has low diagnostic yield and is therefore not profitable in a context where

resources are limited [5]. X-rays of the chest are probably the screening test whose prescription has decreased the most in France helped by the abandonment of its systematic prescription in occupational medicine, obstetrics or preoperatively, the current consensus in France is not to ask for routine radiography under 60 years old [6]. Indeed its reading is difficult and not free of false negatives and especially false positives, resulting in the prescription of additional exams heavy and unnecessary. In some countries, this examination is in great demand by the employees, firstly because of the high proportion of smokers in the workplace who wish to self-control, but also because those workers would like to have a certain advantage at the expense of the employer. For the occupational physicians, the experience of systematic prescriptions for chest X-rays does not show any particular benefit, especially for areas where no potential respiratory risk is to be monitored [6]. Zima-Ebeyard *et al.* in a study entitled Interest of Systematic Diagnostic Radiology of the chest during Periodic Medical Visits (PMV) among workers in Gabon, found a high performance for screening for tuberculosis and low for lung cancer, they therefore propose that thoracic radiography should be performed on a case-by-case basis and should be preceded by a clinical examination including a thorough anamnesis and in particular a good interview on known occupational and environmental exposures [3]. The last similar study carried out in our country dates back to 1998 by Nouthé Djudgang *et al.* who arrived at the same conclusion [7]. Yet in our country the practice of routine chest X-ray in medical periodic visits is still very common, practically requested routinely by almost all working physicians, despite the abandonment by many countries. Thus, we proposed to evaluate the interest of this practice in the follow-up of the workers in our context, in particular as regards the diagnostic yield and the cost-effectiveness ratio.

2. Materials and Methods

It was a cross-sectional descriptive study conducted at the Ngaoundéré Regional Hospital, during a year from January to December 2018, including all persons coming to perform a routine chest X-ray as part of periodic medical visits and who agree to participate in the study, were excluded those who refused to participate in the study and pregnant women for whom X-ray radiographs were contraindicated. A preliminary examination and clinical examination were performed by occupational physicians to search for any underlying illness and the medical history of the person concerned. A chest X-ray with a single posterior-anterior (PA) incidence was then obtained in the radiology and medical imaging department after presenting a request form. The technician after performing the X-ray check the quality before interpretation by a radiologist doctor with a great experience in the interpretation of chest X-rays. The following variables were studied: name, age, sex, workstation, clinical results and chest X-ray results. After interpretation we coded the results of the chest X-ray as normal, sequelae anomaly and evolutionary anomaly depending on the need or not for a comple-

mentary assessment. The results of the chest X-ray were confronted with the clinic and the antecedents. The cost of an X-ray was the one applied in this hospital at the time of the study in local currency which is the *Communauté Financière Africaine* franc (F CFA). This cost was converted into United States dollars (\$) using the exchange rate prevailing at the time of the study. Data were collected with the aid of a standardized questionnaire. The collected data were analyzed statistically using the Sphinx V5 Edition statistical software.

3. Results

A total of 753 workers were enrolled. There were 663 men (88.04%) and 90 women (11.96%), giving a sex ratio of 7:37. The mean age of the study population was 39 years, with a minimum of 26 years and a maximum of 56 years (Figure 1).

Workers were mostly administrative staff with 38.37%, drivers with 17.92%, storekeepers with 15.27% and technicians with 7.43% (Table 1).




Figure 1. Distribution of patients by age.

Table 1. Distribution by job type.

Profession	Frequency	Percentage (%)
Administration	289	38.37
Driver	135	17.92
Storekeeper, handling work	115	15.27
Technician	56	7.43
Accountant	40	5.31
Cashier	33	4.38
Mechanic	1	2.52
Watchman	15	1.99
Others	51	6.77
Total	753	100

An interrogation and clinical examination of the workers before the chest X-ray revealed that 734 (97.47%) were asymptomatic, 19 workers (2.52%) had cough, 16 workers (2.12%) had chest pain, 2 workers (0.26%) dyspnea and one person (0.13%) fever. In addition, there was a history of tuberculosis in 6 workers (0.79%), pleurisy in 8 workers (1.06%) and pulmonary infection in 18 workers (2.39%). 56 cases (7.43%) of occasional smoking and alcohol drinking were declared.

Of the 753 radiographs obtained, 712 were normal (94.55%), while 41 X-rays (5.44%) showed abnormalities (**Table 2**), thus 14 cases (1.85%) of blunting of left costophrenic angle, 9 cases (1.19%) blunting of right costophrenic angle, 8 cases (1.06%) of cardiomegaly, 3 cases (0.39%) of basal infiltrate, 6 cases (0.79%) of apical infiltrate and one case (0.13%) of right pleural liquid effusion of mean abundance with right para-tracheal opacity. Among these abnormalities 40 cases (5.31%) were considered sequelae and one case (0.13%) as evolutionary (**Figure 2**).

The figures below show the examples of sequelae (**Figures 3-5**) and evolutionary (**Figure 6**) abnormalities encountered.

When the results of the chest X-ray were compared with the clinical signs and the pathological history of the workers, it was found that the only case with evolutionary abnormalities on the chest X-ray showed proven clinical signs, 56.52%

Table 2. Distribution of patients by chest X-ray results.

Results	Frequency	Percentage (%)
Normal chest X-ray	712	94.50
Blunting of left costophrenic angle	14	1.80
Blunting of right costophrenic angle	9	1.19
Cardiomegaly	8	1.06
Basal infiltrate	3	0.39
Apical infiltrate	6	0.79
Moderate right pleural effusion + anterior superior right paratracheal opacity	1	0.132
Total	753	100




Figure 2. Distribution of pathology by severity.




Figure 3. Blunting of left costophrenic angle (red arrow).




Figure 4. Right basal infiltrate (red circle).




Figure 5. Moderate right pleural effusion (red arrow), anterior superior right paratracheal opacity (black star) and right hilary opacity (red star).

of the workers with blunting of costophrenic angles on chest X-ray complained of chest pain (**Table 3**) and that one third of those patients had a history of pleurisy, 77.77% of workers with basal or apical infiltrate had a history of either tuberculosis or lung infection (**Table 4**). The cost of a single-view PA chest radiograph




Figure 6. Right apical infiltrate (black star).

Table 3. Distribution of chest X-ray results by clinical signs.

Results	Clinical signs			
	Cough	Chest pain	Dyspnea	fever
Normal chest X-ray	17	02	/	/
Blunting of left and right costo-phrenic angle	/	13	/	/
Cardiomegaly	/	/	01	/
Basal infiltrate	01	/	/	/
Apical infiltrate	/	/	/	/
Moderate right pleural effusion + anterior superior right paratracheal opacity	01	01	01	01

Table 4. Distribution of chest X-ray results by antecedents.

Results	Antecedent		
	Tuberculosis	Pulmonary infection	Pleurisy
Normal chest X-ray	01	10	01
Blunting of left and right costo-phrenic angle	/	02	06
Cardiomegaly	/	/	/
Basal infiltrate	01	01	/
Apical infiltrate	04	01	/
Moderate right pleural effusion + anterior superior right paratracheal opacity	/	01	01

at the Ngaoundere Regional Hospital at the time of the study was 5000 F CFA (\$8.59). The direct cost of all the 753 radiographs obtained was estimated at 3,765,000 F CFA (\$6,469).

4. Discussion

Most of the workers in our studied population were men, this was due to the fact that most of the personnel of the companies occupy posts dedicated to men, such as storekeepers, drivers, assistant drivers and mechanics, posts dedicated to few women, hence the low representation of women in this study. With an average age of 39 years, we think that this represents perfectly the majority of people of working age, able to exercise the different trades found in the companies concerned by these periodic medical visits, the majority of these staff was between 30 and 47 years old, about 78%, Adeko *et al.* [4] in a study of routine chest X-ray during pre-employment medical visits to health workers found an average age of 26.6 years and from 20 to 49 years was predominant, this difference is due to the fact that in this study the people concerned are still looking for work whereas in our study, they are confirmed workers in different companies. In our series, 712 radiographs were normal or 94.55% against 41 cases of abnormalities found or 5.44%, these results are similar to those of Adeko *et al.* who had found in a study in pre-employment 95.2% normal radiographs and 4.8% abnormalities [4], and quite apart from those of Moifo *et al.* who found 2.5% of pathologies in a study evaluating the role of routine chest radiography in students during university enrollment in an area endemic for tuberculosis [5].

Among the abnormalities found in our series, one was evolutionary or 0.13%, in a patient with known and proven clinical manifestations, 40 pathologies or 5.31% were sequelae, as in most studies [5] [7] [8] [9]. Zina-Ebeyard *et al.* in Gabon found 1.5% progressive pathologies in a study of 3276 chest X-rays during periodic medical visits [3]. When the results of the chest X-ray were compared with the clinical signs and the pathological history of the workers, it was found that the only case with evolutionary abnormalities on the chest X-ray showed proven clinical signs, 56.52% of the workers with blunting costo-phrenic angle on chest X-ray complained of chest pain and that one third of these patients had a history of pleurisy, 77.77% of workers with basal or apical infiltrate had a history of either tuberculosis or pulmonary infection. The medical history and clinical signs for which abnormalities were found on the chest X-ray were history of TB and pleurisy. Some abnormalities on chest X-ray were found in the context of chest pain, dyspnea and fever. For this reason, most authors recommend that chest X-rays should be performed on a case-by-case basis after a clinical examination including a thorough anamnesis and in particular a good interrogation on the known occupational and environmental exposures [3]-[10]. Indeed, these chest X-rays were performed in these apparently healthy patients to detect any occult or occupational disease, such as lung cancer, as soon as possible, so that cure is possible, even if the chest X-ray has been shown to have

poor sensitivity for the detection of certain diseases, in particular lung cancer and certain infections, consistent with some studies that have shown that screening is not beneficial even in high-risk groups, in addition, normal radiography does not exclude certain infections such as tuberculosis, which is seen in about 15% of people with normal chest X-rays [11] [12] [13] [14]. Cardiomegaly can be physiological in some people, especially those involved in highly competitive sporting activities.

When exposed to ionizing radiation, the advantages to potential risks should always be carefully considered. If the possible benefits are offset by the potential risks, the procedure can also be considered [5]. We must carefully consider the cost-effectiveness of any diagnostic radiology procedure, particularly in a developing country such as ours, as the expense of resources in procedures with low diagnostic performance would not be cost-effective. The cost of X-rays varies from one country to another, within a country, from one city to another, whatever the cost, the X-ray of the chest must be able to provide a benefit to the patient. In our series, only some patients were able to really benefit from her X-ray, one could really do without X-rays of the chest in other staff.

Our study has some limitations including such as performing a single-incidence chest X-ray, lack of complementary investigations and follow-up of workers with an abnormality, certain conditions such as tuberculosis or cancer could not therefore be excluded in these people.

5. Conclusion

The 5.44% abnormalities rate found in the study appears to be low and calls into question the usefulness and cost-effectiveness of thoracic chest X-ray in periodic medical visits. The only evolutionary pathology was found in a person with proven and known clinical signs. However, patients with lesions on the chest X-ray had clinical signs and antecedents related to these abnormalities, hence the interest of an interrogation and a preliminary clinical examination that could make it possible to discuss the appropriateness of the whether or not the chest X-ray is performed. It should be unsystematic and case-by-case depending on the clinical context and worker history.

Ethical Considerations

This work was done as part of a master's thesis and we had obtained permission from the university and the hospital, the informed consent of the workers, the anonymity was respected in the context of data collection.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References


- [1] Ministry of Labor and Social Security of the Republic of Cameroon. Law N°92/007

of August 14th, 1992 Bearing the Labor Code Decree N°79/096 of March 21st, 1979 Fixing the Modalities of Exercise of the Occupational Medicine.

<https://www.camerlex.com/cameroun-l-organisation-du-ministere-du-travail-et-de-la-securite-sociale-20647/>

- [2] College of University Hospital Teachers of Occupational Medicine of Ile-de-France and Labor Inspection of Ile de France. Legislation and Regulations in Occupational Health. January 2006. <http://www.uvmt.org/Multimedia/Poly20052006.pdf>
- [3] Zima-Ebeyard, M.M., Djeki, J., Ondo, Y.S., Nambo-Wezet, G.B. and Obiang Ossoubita, B. (2009) Interest of the Practice of Systematic Chest Radiography during Periodic Medical Visits of Workers in Gabon. *CAMIP* 2009, January-March 2009. <http://www.camip.info/Discussion,624.html>
- [4] Adeko, O.O., Ariba, A.J., Olatunji, A.A. and Toyobo, O.O. (2017) Routine Chest Radiograph in Pre-Employment Medical Examination for Healthcare Workers: Time for a Review of the Protocol. *Nigerian Postgraduate Medical Journal*, **24**, 93-96.
- [5] Boniface, M., Joshua, T., Walter, P.E., Fernande, Z.O., Emrick, G.K. and Joseph, G.F. (2012) Assessing the Role of Routine Chest Radiography in Asymptomatic Students during Registration at a University in an Endemic Area of Tuberculosis. *Annals of Tropical Medicine and Public Health*, **5**, 419-422. <https://doi.org/10.4103/1755-6783.105122>
- [6] Doctravail (2011) X-Ray of the Non-Systematic Lungs. The Doctor's Gazette 19 April 2011. <http://medecinetravail.canalblog.com/archives/2011/04/19/20931591.html>
- [7] Nouthé Djubgang, J. and Fayomi, E.B. (1998) Interest of the Practice of the Systematic Pulmonary X-Ray during the Periodic Medical Visits of the Workers in Cameroon. *Revue de Médecine du Travail*, **25**, 95-98.
- [8] Jessica, V. (2014) Interest of the Complementary Examinations Carried out and Prescribed during the Systematic Medical Visits. *Human Medicine and Pathology*. Articles L335.2-L335.10. <http://www.dumas.ccsd.cnrs.fr/>
- [9] Malnick, S., Duek, G., Beilinson, N., Neogolani, V., Basevitz, A., Somin, M., Cohen, J., Katz, M. and Schattner, A. (2010) Routine Chest X-Ray on Hospital Admission: Does It Contribute to Diagnosis or Treatment? *The Israel Medical Association Journal*, **12**, 357-361.
- [10] Chukwuani, A.E., Osanaiye, A., Eneje, O.F. and Omiyi, D.O. (2017) An Audit of Routine Chest X-Ray Done During Pre-Employment Medical Screening in South-Western Nigeria. *Central African Journal of Public Health*, **3**, 51-54.
- [11] Shah, P.K., Astin, J.H., White, C.S., Patel, P., Haramati, L.B., Pearson, G.D., *et al.* (2003) Missed Non-Small Cell Lung Cancer: Radiographic Findings of Potentially Resectable Lesions Evident Only in Retrospect. *Radiology*, **226**, 235-241. <https://doi.org/10.1148/radiol.2261011924>
- [12] Fontana, R.S., Sanderson, D.R., Woolner, L.B., Taylor, W.F., Miller, W.E. and Muhm, J.R. (1986) Lung Cancer Screening: The Mayo Program. *Journal of Occupational Medicine*, **28**, 746-750.
- [13] Marcus, P.M., Bergstralh, E.J., Fagerstrom, R.M., Williams, D.E., Fontana, R., Taylor, W.F. and Prorok, P.C. (2000) Lung Cancer Mortality in the Mayo Lung Project: Impact of Extended Follow-Up. *Journal of the National Cancer Institute*, **92**, 1308-1316. <https://doi.org/10.1093/jnci/92.16.1308>
- [14] Woodring, J.H., Vandiviere, H.M., Fried, A.M., Dillon, M.L., Williams, T.D. and Melvin, I.G. (1986) Update: The Radiographic Features of Pulmonary Tuberculosis. *American Journal of Roentgenology*, **146**, 497-506. <https://doi.org/10.2214/ajr.146.3.497>

Call for Papers



Advances in Molecular Imaging (AMI)

ISSN 2161-6728 (Print) ISSN 2161-6752 (Online)

<http://www.scirp.org/journal/ami>

Advances in Molecular Imaging is a peer-reviewed, open access journal that publishes original research articles, review articles, case reports, and clinical studies in all areas of molecular imaging:

Editor-in-Chief

Prof. Orhan Nalcioglu

University of California-Irvine, USA

Editorial Board

Prof. Gjumrakch Aliev
Dr. Ying Bai
Prof. Baowei Fei
Dr. Zhong-Ping Feng
Prof. Richard Hans Gomer
Prof. Gultekin Gulsen
Prof. Mohammad Mojammel Al Hakim
Dr. Maria Kempe
Dr. David Sigmund Liebeskind
Dr. Stefan Lorkowski
Prof. Kenneth Maiese
Prof. Adalberto Merighi

Prof. Jean-Pierre Raufman
Prof. Gianfranco Risuleo
Prof. Steven Alan Rosenzweig
Prof. Phillip Ruiz
Dr. Joy Sinha
Prof. Maurizio Sorice
Prof. Lun-Quan Sun
Prof. Bin Tean Teh
Prof. Masakazu Toi
Prof. Horst Christian Weber
Dr. Jia Lin Yang

Subject Coverage

Chemical Synthesis Platform Technologies
Computer Vision and Image Understanding
Contrast Media & Molecular Imaging
Imaging & Microscopy
Imaging Decisions MRI
Imaging Systems and Technology
ImmunoPET and ImmunoSPECT
In Vitro Molecular Diagnostics (IVMD)
Molecular Diagnostics in Cancer and Immune Disorders

Molecular Imaging and Biology
Molecular Imaging Using ImmunoPET
MRI/MDCT/PET Imaging of Various Organs
Neuroimaging
Nuclear Medicine and Molecular Imaging
Onco-Imaging
Preclinical Imaging Systems
Systems Biology
Tracer and Pharmacokinetic Modeling

Notes for Intending Authors

We are also interested in: 1) Short reports—2-5 page papers in which an author can either present an idea with a theoretical background but has not yet completed the research needed for a complete paper or preliminary data; 2) Book reviews—Comments and critiques.

Website and E-Mail

<http://www.scirp.org/journal/ami>

E-mail: ami@scirp.org

What is SCIRP?

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

What is Open Access?

All original research papers published by SCIRP are made freely and permanently accessible online immediately upon publication. To be able to provide open access journals, SCIRP defrays operation costs from authors and subscription charges only for its printed version. Open access publishing allows an immediate, worldwide, barrier-free, open access to the full text of research papers, which is in the best interests of the scientific community.

- High visibility for maximum global exposure with open access publishing model
- Rigorous peer review of research papers
- Prompt faster publication with less cost
- Guaranteed targeted, multidisciplinary audience



**Scientific
Research
Publishing**

Website: <http://www.scirp.org>

Subscription: sub@scirp.org

Advertisement: service@scirp.org