

Ground States for a Class of Nonlinear Schrodinger-Poisson Systems with Positive Potential

Guoqing Zhang*, Xue Chen

College of Sciences, University of Shanghai for Science and Technology, Shanghai, China
Email: shzhangguoqing@126.com

Received 10 November 2014; revised 29 November 2014; accepted 15 December 2014

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Based on Nehari manifold, Schwarz symmetric methods and critical point theory, we prove the existence of positive radial ground states for a class of Schrodinger-Poisson systems in \mathbb{R}^3 , which doesn't require any symmetry assumptions on all potentials. In particular, the positive potential is interesting in physical applications.

Keywords

Ground States, Schrodinger-Poisson Systems

1. Introduction

In this paper, we consider the following nonlinear Schrodinger-Poisson systems

$$\begin{cases} -\Delta u + V(x)u - \lambda \rho(x)\Phi u + Q(x)|u|^{p-2}u = 0, & x \in \mathbb{R}^3, \\ -\Delta \Phi = \rho(x)u^2, & x \in \mathbb{R}^3, \end{cases} \quad (1.1)$$

where $\lambda > 0$, $2 < p < 4$; $V(x)$, $\rho(x)$ and $Q(x)$ are positive potentials defined in \mathbb{R}^3 .

In recent years, such systems have been paid great attention by many authors concerning existence, non-existence, multiplicity and qualitative behavior. The systems are to describe the interaction of nonlinear Schrodinger field with an electromagnetic field. When $\lambda = -1$, $V(x) = \rho(x) = 1$, $Q(x) = -1$, the existence of non-trivial solution for the problem (1.1) was proved as $p \in (4, 6)$ in [1], and non-existence result for $p \in (0, 2]$ or $p \in (6, +\infty)$ was proved in [2]. When $\lambda < 0$, $V(x) = \rho(x) = 1$, $Q(x) = -1$, using critical point theory,

*Corresponding author.

Ruiz [3] obtained some multiplicity results for $p \in (2, 3)$, and existence results for $p \in [3, 6)$. Later, Ambrosetti and Ruiz [4], and Ambrosetti [5] generalized some existence results of Ruiz [3], and obtained the existence of infinitely solutions for the problem (1.1).

In particular, Sanchel and Soler [6] considered the following Schrodinger-Poisson-Slater systems

$$\begin{cases} -\Delta u + \omega u + \Phi u - |u|^{\frac{2}{3}} u = 0, & x \in \mathbb{R}^3, \\ -\Delta \Phi = u^2, & x \in \mathbb{R}^3, \end{cases} \quad (1.2)$$

where $\omega \in \mathbb{R}$. The problem (1.2) was introduced as the model of the Hartree-Fock theory for a one-component plasma. The solution is obtained by using the minimization argument and ω as a Lagrange multiplier. However, it is not known if the solution for the problem (1.2) is radial. Mugani [7] considered the following generalized Schrodinger-Poisson systems

$$\begin{cases} -\Delta u + \omega u - \lambda \Phi u + W_u(x, u) = 0, & x \in \mathbb{R}^3, \\ -\Delta \Phi = u^2, & x \in \mathbb{R}^3, \end{cases} \quad (1.3)$$

where $\omega \in \mathbb{R}$, $\lambda > 0$ and $W(x, s) = W(|x|, s)$, and proved the existence of radially symmetric solitary waves for the problem (1.3).

In this paper, without requiring any symmetry assumptions on $V(x)$, $\rho(x)$ and $Q(x)$, we obtain the existence of positive radial ground state solution for the problem (1.1). In particular, the positive potential $Q(x)$ implies that we are dealing with systems of particles having positive mass. It is interesting in physical applications.

The paper is organized as following. In Section 2, we collect some results and state our main result. In Section 3, we prove some lemmas and consider the problem (1.1) at infinity. Section 4 is devoted to our main theorem.

2. Preliminaries and Main Results

Let $L^s(\mathbb{R}^3)$, $1 \leq s < +\infty$ denotes a Lebesgue space, the norm in $L^s(\mathbb{R}^3)$ is $\|u\|_s = \left(\int_{\mathbb{R}^3} |u|^s dx \right)^{\frac{1}{s}}$, $D^{1,2}(\mathbb{R}^3)$ is the completion of $C_0^\infty(\mathbb{R}^3)$ with respect to the norm

$$\|u\|_{D^{1,2}} = \left(\int_{\mathbb{R}^3} |\nabla u|^2 dx \right)^{\frac{1}{2}}$$

$H^1(\mathbb{R}^3)$ be the usual Sobolev space with the usual norm

$$\|u\|_{H^1} = \left(\int_{\mathbb{R}^3} (|\nabla u|^2 + |u|^2) dx \right)^{\frac{1}{2}}.$$

Assume that the potential $V(x)$ satisfies

H1) $V(x) \in C^1(\mathbb{R}^3, \mathbb{R})$, $\inf_{x \in \mathbb{R}^3} V(x) = 1$, $V(x) \leq V_\infty = \lim_{|x| \rightarrow \infty} V(x) < \infty$.

Let $H_V^1(\mathbb{R}^3)$ be the Hilbert subspace of $u \in H^1(\mathbb{R}^3)$ such that

$$\|u\|_{H_V^1} = \left(\int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)|u|^2) dx \right)^{\frac{1}{2}} \quad (2.1)$$

Then $H_V^1(\mathbb{R}^3) \subset H^1(\mathbb{R}^3) \subset L^s(\mathbb{R}^3)$, $2 \leq s \leq 6$ with the corresponding embeddings being continuous (see [8]). Furthermore, assume the potential $\rho(x)$ satisfies

H2) $\rho(x) > 0$, $\lim_{|x| \rightarrow +\infty} \rho(x) = \rho_\infty > 0$, $\rho_0(x) = \rho(x) - \rho_\infty \in L^2(\mathbb{R}^3)$.

It is easy to reduce the problem (1.1) to a single equation with a non-local term. Indeed, for every $v \in D^{1,2}(\mathbb{R}^3)$, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^3} \rho(x) u^2 v dx \right| &= \left| \int_{\mathbb{R}^3} (\rho(x) - \rho_\infty) u^2 v dx + \int_{\mathbb{R}^3} \rho_\infty u^2 v dx \right| \\ &\leq \int_{\mathbb{R}^3} |\rho(x) - \rho_\infty| u^2 |v| dx + \int_{\mathbb{R}^3} \rho_\infty u^2 |v| dx \\ &\leq \left(\int_{\mathbb{R}^3} (|\rho_0(x)| u^2)^{\frac{6}{5}} dx \right)^{\frac{5}{6}} \left(\int_{\mathbb{R}^3} (v)^6 dx \right)^{\frac{1}{6}} + \rho_\infty \left(\int_{\mathbb{R}^3} (u^2)^{\frac{6}{5}} dx \right)^{\frac{5}{6}} \left(\int_{\mathbb{R}^3} (v)^6 dx \right)^{\frac{1}{6}} \quad (2.2) \\ &\leq |v|_6 \left[|\rho_0(x)|_2 \left(\int_{\mathbb{R}^3} (u^2)^3 dx \right)^{\frac{1}{3}} + \rho_\infty \left(\int_{\mathbb{R}^3} (u^2)^{\frac{6}{5}} dx \right)^{\frac{5}{6}} \right]. \end{aligned}$$

Since $\rho_0(x) = \rho(x) - \rho_\infty \in L^2(\mathbb{R}^3)$, $u \in H^1(\mathbb{R}^3)$ and (2.1), by the Lax-Milgram theorem, there exists a unique $\Phi[u]$ such that

$$\int_{\mathbb{R}^3} \nabla \Phi[u] \nabla v dx = \int_{\mathbb{R}^3} \rho(x) u^2 v dx, \quad \forall v \in D^{1,2}(\mathbb{R}^3) \quad (2.3)$$

It follows that $\Phi[u]$ satisfies the Poisson equation

$$-\Delta \Phi[u] = \rho(x) u^2$$

and there holds

$$-\Phi[u] = \int_{\mathbb{R}^3} \frac{\rho(x) u^2(y)}{|x-y|} dy = \frac{1}{|x|} * \rho(x) u^2$$

Because $\rho(x) > 0$, we have $\Phi[u] > 0$ when $u \neq 0$, and $\|\Phi[u]\|_{D^{1,2}} = M \|u\|_{H^1}^2$, M is positive constant. Substituting $\Phi[u]$ in to the problem (1.1), we are lead to the equation with a non-local term

$$-\Delta u + V(x)u - \lambda \rho(x) \Phi[u] u + Q(x) |u|^{p-2} u = 0. \quad (2.4)$$

In the following, we collect some properties of the functional $\Phi[u]$, which are useful to study our problem.

Lemma 2.1. [9] For any $u \in H^1(\mathbb{R}^3)$, we have

- 1) $\Phi[u]: H^1(\mathbb{R}^3) \rightarrow D^{1,2}(\mathbb{R}^3)$ is continuous, and maps bounded sets into bounded sets;
- 2) if $u_n \rightarrow u$ weakly in $H^1(\mathbb{R}^3)$, then $\Phi[u_n] \rightarrow \Phi[u]$ weakly in $D^{1,2}(\mathbb{R}^3)$;
- 3) $\Phi[tu(x)] = t^2 \Phi[u(x)]$ for all $t \in \mathbb{R}$.

Now, we state our main theorem in this paper.

Theorem 2.2. Assume that $\lambda > 0$, $2 < p < 4$, the potential $V(x)$ satisfies condition H1), the potential $\rho(x)$ satisfies condition H3) and $\rho(x) \geq \rho_\infty$, the potential $Q(x)$ satisfies

$$H3) \quad Q(x) > 0, \quad \lim_{|x| \rightarrow +\infty} Q(x) = Q_\infty > 0, \quad Q_0(x) = Q(x) - Q_\infty \in L^{\frac{6}{6-p}}(\mathbb{R}^3)$$

and $Q(x) \leq Q_\infty$, $Q(x) - Q_\infty < 0$ on positive measure. Then there exists a positive radial ground state solution for the problem (1.1).

Remark 2.3. If $\lambda \leq 0$, $V(x)$, $\rho(x)$ and $Q(x)$ are positive potentials defined in \mathbb{R}^3 , and $2 < p < 6$, $(u, \Phi) \in H_V^1(\mathbb{R}^3) \times D^{1,2}(\mathbb{R}^3)$ be a solution for the problem (1.1). Then $(u, \Phi) = (0, 0)$, Indeed, we have

$$\begin{aligned} 0 &= \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2) dx - \lambda \int_{\mathbb{R}^3} \rho(x) \Phi[u] u^2 dx + \int_{\mathbb{R}^3} Q(x) |u|^p dx \\ &\geq \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2) dx. \end{aligned}$$

Since $V(x) > 0$, this implies $u = 0$. By Lemma 2.1, we have $\Phi = 0$.

3. Some Lemmas and the Problem (1.1) at Infinity

Now, we consider the functional $I_\lambda : H_V^1(\mathbb{R}^3) \rightarrow \mathbb{R}$ given by

$$\begin{aligned} I_\lambda(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 + V(x)u^2) dx - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx + \frac{1}{p} \int_{\mathbb{R}^3} Q(x)|u|^p dx \\ &= \frac{1}{2} \|u\|_{H_V^1}^2 - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx + \frac{1}{p} \int_{\mathbb{R}^3} Q(x)|u|^p dx. \end{aligned}$$

Since $\rho(x)$ satisfies condition H2), by (2.2), the Holder inequality and Sobolev inequality, we have

$$\int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx \leq \bar{S}^{-1} \left[S_6^{-2} |\rho|_2 + \rho_\infty S_{\left(\frac{12}{5}\right)}^{-2} \right] \|u\|_{H_V^1}^2 \|v\|_{D^{1,2}}, \quad (3.2)$$

where $\bar{S} = \inf_{u \in D^{1,2}(\mathbb{R}^3) \setminus \{0\}} \frac{\|u\|_{D^{1,2}}}{|u|_6}$ and $S = \inf_{u \in H_V^1(\mathbb{R}^3) \setminus \{0\}} \frac{\|u\|_{H_V^1}}{|u|_6}$. Since the potential $Q(x)$ satisfies condition Q, $2 < p < 4$, we have

$$\begin{aligned} \int_{\mathbb{R}^3} Q(x)|u|^p dx &\leq \left| \int_{\mathbb{R}^3} (Q(x) - Q_\infty)|u|^p dx + \int_{\mathbb{R}^3} Q_\infty |u|^p dx \right| \\ &\leq \int_{\mathbb{R}^3} |Q_0(x)| |u|^p dx + \int_{\mathbb{R}^3} Q_\infty |u|^p dx \\ &\leq \left(\int_{\mathbb{R}^3} |Q_0(x)|^{\left(\frac{6}{6-p}\right)} dx \right)^{\left(\frac{6-p}{6}\right)} \left(\int_{\mathbb{R}^3} (u)^6 dx \right)^{\left(\frac{p}{6}\right)} + Q_\infty \int_{\mathbb{R}^3} |u|^p dx. \end{aligned}$$

By Sobolev inequality, we obtain that

$$\int_{\mathbb{R}^3} Q(x)|u|^p dx \leq M \|u\|_{H_V^1}^p \quad (3.3)$$

Combining (3.2) and (3.3), we obtain that the functional I_λ is a well defined C^1 functional, and if $u \in H_V^1(\mathbb{R}^3)$ is critical point of it, then the pair $(u, \Phi[u])$ is a weak solution of the problem (1.1).

Now, we define the Nehari manifold ([10]) of the functional I_λ

$$N_\lambda = \{u \in H_V^1(\mathbb{R}^3) \setminus \{0\} : H_\lambda(u) = 0\},$$

where

$$H_\lambda(u) = I'_\lambda(u)[u] = \|u\|_{H_V^1}^2 - \lambda \int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx + \int_{\mathbb{R}^3} Q(x)|u|^p dx$$

Hence, we have

$$\begin{aligned} I'_\lambda(u)|_{N_\lambda} &= \frac{1}{4} \|u\|_{H_V^1}^2 + \left(\frac{1}{p} - \frac{1}{4}\right) \int_{\mathbb{R}^3} Q(x)|u|^p dx \\ &= \left(\frac{1}{2} - \frac{1}{p}\right) \|u\|_{H_V^1}^2 + \lambda \left(\frac{1}{p} - \frac{1}{4}\right) \int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx \\ &= \left(\frac{1}{2} - \frac{1}{p}\right) \int_{\mathbb{R}^3} Q(x)|u|^p dx - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho(x)\Phi[u]u^2 dx. \end{aligned} \quad (3.4)$$

Lemma 3.1. 1) For any $\lambda > 0$, $u \in H_V^1(\mathbb{R}^3) \setminus \{0\}$, there exists a unique $t(u) > 0$ such that $t(u)u \in N_\lambda$. Moreover, we have $I_\lambda(t(u)u) = \max_{t \geq 0} I_\lambda(tu)$.

2) $I_\lambda(u)$ is bounded from below on N_λ by a positive solution.

Proof. 1) Taking any $u \in H_V^1(\mathbb{R}^3) \setminus \{0\}$ and $\|u\|_{H_V^1} = 1$, we obtain that there exists a unique $t(u) > 0$ such that $t(u)u \in N_\lambda$. Indeed, we define the function $g(t) = I_\lambda(tu)$. We note that $g'(t) = (I'_\lambda(tu), v) = 0$ if only if $tu \in N_\lambda$. Since $g'(t) = 0$ is equivalent to

$$t^2 \|u\|_{H_V^1}^2 - \lambda t^4 \int_{\mathbb{R}^3} \rho(x) \Phi[u] u^2 dx + t^p \int_{\mathbb{R}^3} Q(x) |u|^p dx = 0.$$

By $\rho(x), Q(x) > 0$ and $\Phi[u] > 0$, we have

$$b = \int_{\mathbb{R}^3} \rho(x) \Phi[u] u^2 dx > 0, \quad c = \int_{\mathbb{R}^3} Q(x) |u|^p dx > 0.$$

By $\lambda > 0$, $2 < p < 4$, the equation $1 - \lambda b t^2 + c t^{p-2} = 0$ has a unique $t(u) > 0$ and the corresponding point $t(u)u \in N_\lambda$ and $I_\lambda(t(u)u) = \max_{t \geq 0} I_\lambda(tu)$.

2) Let $u \in N_\lambda$, by (3.4) and $2 < p < 4$, we have

$$\begin{aligned} I_\lambda(u) &= \left(\frac{1}{2} - \frac{1}{p} \right) \|u\|_{H_V^1}^2 + \lambda \left(\frac{1}{p} - \frac{1}{4} \right) \int_{\mathbb{R}^3} \rho(x) \Phi[u] u^2 dx \\ &\geq \left(\frac{1}{2} - \frac{1}{p} \right) \|u\|_{H_V^1}^2 > C > 0. \end{aligned}$$

By the definition of Nehari manifold N_λ of the functional I_λ , we obtain that u is a critical point of I_λ if and only if u is a critical point of I_λ constrained on N_λ . (3.5)

Now, we set

$$m_\lambda = \inf \{ I_\lambda(u) : u \in N_\lambda \}$$

By 2) of Lemma 3.1, we have $m_\lambda > 0$.

Since $\lim_{|x| \rightarrow \infty} V(x) = V_\infty$, $\lim_{|x| \rightarrow +\infty} \rho(x) = \rho_\infty$, $\lim_{|x| \rightarrow +\infty} Q(x) = Q_\infty$, we consider the problem (1.1) at infinity

$$\begin{cases} -\Delta u + V_\infty u - \lambda \rho_\infty \Phi u + Q_\infty |u|^{p-2} u = 0, & x \in \mathbb{R}^3, \\ -\Delta \Phi = \rho_\infty u^2, & x \in \mathbb{R}^3. \end{cases} \quad (3.6)$$

Similar to (2.2), we obtain that there exists a unique $\tilde{\Phi}[u]$ such that

$$\int_{\mathbb{R}^3} \nabla \tilde{\Phi}[u] \nabla v dx = \int_{\mathbb{R}^3} \rho_\infty u^2 v dx, \quad \forall v \in D^{1,2}(\mathbb{R}^3).$$

It follows that $\tilde{\Phi}[u]$ satisfies the Poisson equation

$$-\Delta \tilde{\Phi}[u] = \rho_\infty u^2 \quad (3.7)$$

Hence substituting $\tilde{\Phi}[u]$ into the first equation of (3.6) we have to study the equivalent problem

$$-\Delta u + V_\infty u - \lambda \rho_\infty \tilde{\Phi}[u] u + Q_\infty |u|^{p-2} u = 0 \quad (3.8)$$

The weak solution of the problem (3.8) is the critical point of the functional

$$\begin{aligned} I_\lambda^\infty(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla u|^2 + V_\infty u^2) dx - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u] u^2 dx + \frac{1}{p} \int_{\mathbb{R}^3} Q_\infty |u|^p dx \\ &= \frac{1}{2} \|u\|_{H_{V_\infty}^1}^2 - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u] u^2 dx + \frac{1}{p} \int_{\mathbb{R}^3} Q_\infty |u|^p dx, \end{aligned}$$

where $H_{V_\infty}^1(\mathbb{R}^3) = H^1(\mathbb{R}^3)$ is endowed with the norm

$$\|u\|_{H_{V_\infty}^1} = \left(\int_{\mathbb{R}^3} (|\nabla u|^2 + V_\infty u^2) dx \right)^{\frac{1}{2}}$$

Define the Nehari manifold of the functional I_λ^∞

$$N_\lambda^\infty = \{u \in H_{V_\infty}^1(\mathbb{R}^3) \setminus \{0\} : H_\lambda^\infty(u) = 0\},$$

where

$$H_\lambda^\infty(u) = I_\lambda^{\prime\infty}(u)[u] = \|u\|_{H_{V_\infty}^1}^2 - \lambda \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u] u^2 dx + \int_{\mathbb{R}^3} Q_\infty |u|^p dx$$

and

$$m_\lambda^\infty = \inf \{I_\lambda^\infty(u) : u \in N_\lambda^\infty\} > 0$$

The Nehari manifold N_λ^∞ has properties similar to those of N_λ .

Lemma 3.2. The problem (3.8) has a positive radial ground state solution $\omega_\infty \in N_\lambda^\infty$ such that

$$I_\lambda^\infty(\omega_\infty) = m_\lambda^\infty$$

For the proof of Lemma 3.2, we make use of Schwarz symmetric method. We begin by recalling some basic properties.

Let $f \in L^s(\mathbb{R}^3)$ such that $f \geq 0$, then there is a unique nonnegative function $f^* \in L^s(\mathbb{R}^3)$, called the Schwarz symmetric of f , such that it depends only on $|x|$, whose level sets

$$\{x \in \mathbb{R}^3 : f(x) > t\} = \{x \in \mathbb{R}^3 : f^*(x) > t\}.$$

We consider the following Poisson equation

$$-\Delta \phi = f \quad \text{and} \quad -\Delta v = f^*$$

From Theorem 1 of [11], we have

$$\int_{\mathbb{R}^3} |\nabla v|^s dx \geq \int_{\mathbb{R}^3} |\nabla \phi|^s dx, \quad \forall 0 < s \leq 2.$$

Hence, let $\phi = \tilde{\Phi}[u]$, $f = \rho_\infty u^2$ and $v = \tilde{\Phi}[u^*]$, $f^* = \rho_\infty (u^*)^2$, we have

$$\int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u] u^2 dx \leq \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u^*] (u^*)^2 dx. \quad (3.9)$$

The Proof of Lemma 3.2. Let $u_n \in N_\lambda^\infty$ be such that $I_\lambda^\infty(u_n) \rightarrow m_\lambda^\infty$. Let $t_n > 0$ such that $t_n |u_n| \in N_\lambda^\infty$ then we have

$$\int_{\mathbb{R}^3} (|\nabla u_n|^2 + V_\infty (u_n)^2) dx - \lambda \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u_n] (u_n)^2 dx + \int_{\mathbb{R}^3} Q_\infty |u_n|^p dx = 0,$$

and

$$(t_n)^2 \int_{\mathbb{R}^3} (|\nabla u_n|^2 + V_\infty (u_n)^2) dx - \lambda (t_n)^4 \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[u_n] (u_n)^2 dx + (t_n)^p \int_{\mathbb{R}^3} Q_\infty |u_n|^p dx = 0.$$

Hence, we obtain that

$$\left((t_n)^2 - (t_n)^4 \right) \int_{\mathbb{R}^3} (|\nabla u_n|^2 + V_\infty (u_n)^2) dx + \left((t_n)^p - (t_n)^4 \right) \int_{\mathbb{R}^3} Q_\infty |u_n|^p dx = 0. \quad (3.10)$$

Since $t_n > 0$ and $2 < p < 4$, (3.10) implies that $t_n = 1$. Therefore, we can assume that $u_n \geq 0$. On the other hand, let $(u_n)^*$ be the Schwartz symmetric function associated to u_n , then we have

$$\int_{\mathbb{R}^3} |u_n|^p dx = \int_{\mathbb{R}^3} |(u_n)^*|^p dx, \quad \text{and} \quad \int_{\mathbb{R}^3} |\nabla u_n|^2 dx \geq \int_{\mathbb{R}^3} |\nabla (u_n)^*|^2 dx \quad (3.11)$$

Let $(t_n)^* > 0$ be such that $(t_n)^* (u_n)^* \in N_\lambda^\infty$, and $u_n \in N_\lambda^\infty$, by (3.9) and (3.11), we have

$$\begin{aligned} 0 &= \left[(t_n)^* \right]^2 \int_{\mathbb{R}^3} \left(|\nabla (u_n)^*|^2 + V_\infty \left[(u_n)^* \right]^2 \right) dx - \lambda \left[(t_n)^* \right]^4 \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi} \left[(u_n)^* \right] \left[(u_n)^* \right]^2 dx + \left[(t_n)^* \right]^p \int_{\mathbb{R}^3} Q_\infty \left| (u_n)^* \right|^p dx \\ &\leq \left[(t_n)^* \right]^2 \int_{\mathbb{R}^3} \left(|\nabla u_n|^2 + V_\infty (u_n)^2 \right) dx - \lambda \left[(t_n)^* \right]^4 \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi} [u_n] (u_n)^2 dx + \left[(t_n)^* \right]^p \int_{\mathbb{R}^3} Q_\infty |u_n|^p dx \\ &= \left\{ \left[(t_n)^* \right]^2 - \left[(t_n)^* \right]^4 \right\} \int_{\mathbb{R}^3} \left(|\nabla u_n|^2 + V_\infty (u_n)^2 \right) dx + \left\{ \left[(t_n)^* \right]^p - \left[(t_n)^* \right]^4 \right\} \int_{\mathbb{R}^3} Q_\infty |u_n|^p dx. \end{aligned}$$

This implies that $(t_n)^* \leq 1$. Therefore, we have $I_\lambda^\infty \left[(u_n)^* \right] \leq I_\lambda^\infty (u_n)$, and we can suppose that u_n is radial in $H_{V_\infty}^1(\mathbb{R}^3)$. Since $H_{V_\infty, r}^1(\mathbb{R}^3)$ is compactly embedded into $L^p(\mathbb{R}^3)$ for $2 < p < 4$, we obtain that m_λ^∞ is achieved at some $\omega_\infty \in N_\lambda^\infty$ which is positive and radial. Therefore, Lemma 3.2 is proved.

4. The Proof of Main Theorem

In this section, we prove Theorem 2.2. Firstly, we consider a compactness result and obtain the behavior of the (PS) sequence of the functional I_λ .

Lemma 4.1. Let u_n be a (PS)_d sequence of the functional I_λ constrained on N_λ , that is

$$u_n \in N_\lambda, \quad I_\lambda(u_n) \rightarrow d \quad \text{and} \quad I'_\lambda(u_n)|_{N_\lambda} \rightarrow 0, \quad \text{as } n \rightarrow \infty \quad (4.1)$$

Then there exists a solution u of the problem (2.4), a number $k \in \mathbb{N} \cup \{0\}$, k functions u^1, u^2, \dots, u^k of $H_V^1(\mathbb{R}^3)$ and k sequences of points y_n^j , $0 \leq j \leq k$ such that

$$1) \quad |y_n^j| \rightarrow +\infty, \quad |y_n^j - y_n^i| \rightarrow +\infty, \quad \text{if } i \neq j, \quad n \rightarrow \infty;$$

$$2) \quad u_n - \sum_{j=1}^k u^j(\cdot - y_n^j) \rightarrow \bar{u};$$

$$3) \quad I_\lambda(u_n) \rightarrow I_\lambda(\bar{u}) + \sum_{j=1}^k I_\lambda^\infty(u^j);$$

$$4) \quad u^j \text{ are non-trivial weak solution of the problem (3.8).}$$

Proof. The proof is similar to that of Lemma 4.1 in [9].

By Lemma 4.1, taking into account that $I_\lambda^\infty(u^j) \geq m_\lambda$ for all j and $d \in (0, m_\lambda)$, we obtain that $k = 0$ and $u_n \rightarrow \bar{u}$ in $H_V^1(\mathbb{R}^3)$ (strongly), i.e. u_n is relatively compact for all $d \in (0, m_\lambda)$. Hence we only need to prove that the energy of a solution of the problem (2.4) cannot overcome the energy of a ground state solution of the problem (3.8).

The proof of Theorem 2.2. By Lemma 4.1, we only prove that $m_\lambda < m_\lambda^\infty$. Indeed, let $\omega_\infty \in N_\lambda^\infty$ such that $I_\lambda^\infty(\omega_\infty) = m_\lambda^\infty$, and let $t > 0$ such that $t\omega_\infty \in N_\lambda$. Since $V(x) \leq V_\infty$, $\rho(x) \geq \rho_\infty$ and $Q(x) \leq Q_\infty$, we have

$$\begin{aligned} m_\lambda &\leq I_\lambda(t\omega_\infty) = \frac{t^2}{2} \int_{\mathbb{R}^3} \left(|\nabla \omega_\infty|^2 + V(x) \omega_\infty^2 \right) dx - \frac{\lambda t^4}{4} \int_{\mathbb{R}^3} \rho(x) \Phi[\omega_\infty] \omega_\infty^2 dx + \frac{t^p}{p} \int_{\mathbb{R}^3} Q(x) |\omega_\infty|^p dx \\ &\leq \frac{t^2}{2} \int_{\mathbb{R}^3} \left(|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2 \right) dx - \frac{\lambda t^4}{4} \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[\omega_\infty] \omega_\infty^2 dx + \frac{t^p}{p} \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx. \end{aligned} \quad (4.2)$$

Since $\omega_\infty \in N_\lambda^\infty$ and $t\omega_\infty \in N_\lambda$, we have

$$\begin{aligned}
t^4 \int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2) dx + t^4 \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx &= \lambda t^4 \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[\omega_\infty] \omega_\infty^2 dx \\
&\leq \lambda t^4 \int_{\mathbb{R}^3} \rho(x) \Phi[\omega_\infty] \omega_\infty^2 dx \\
&= t^2 \int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V(x) \omega_\infty^2) dx + t^p \int_{\mathbb{R}^3} Q(x) |\omega_\infty|^p dx \\
&\leq t^2 \int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2) dx + t^p \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx.
\end{aligned}$$

Therefore, we have

$$(t^4 - t^2) \int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2) dx + (t^4 - t^p) \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx \leq 0$$

By $2 < p < 4$, we have $t \leq 1$. If $t = 1$, we have $\omega_\infty \in N_\lambda^\infty$ and $\omega_\infty \in N_\lambda$. Hence, by $\omega_\infty \in N_\lambda^\infty$, we have

$$\int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2) dx + \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx = \lambda \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[\omega_\infty] \omega_\infty^2 dx \quad (4.3)$$

and by $\omega_\infty \in N_\lambda$, we have

$$\int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V(x) \omega_\infty^2) dx + \int_{\mathbb{R}^3} Q(x) |\omega_\infty|^p dx = \lambda \int_{\mathbb{R}^3} \rho(x) \Phi[\omega_\infty] \omega_\infty^2 dx. \quad (4.4)$$

Combining (4.3) and (4.4), we have

$$\int_{\mathbb{R}^3} (V_\infty - V(x)) \omega_\infty^2 dx + \int_{\mathbb{R}^3} (Q_\infty - Q(x)) |\omega_\infty|^p dx - \lambda \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[\omega_\infty] \omega_\infty^2 dx + \lambda \int_{\mathbb{R}^3} \rho(x) \Phi[\omega_\infty] \omega_\infty^2 dx = 0$$

Since $V(x) \leq V_\infty$, $\rho(x) \geq \rho_\infty$, $Q(x) \leq Q_\infty$, and $Q(x) - Q_\infty < 0$ on a positive measure, we have

$$\int_{\mathbb{R}^3} (Q_\infty - Q(x)) |\omega_\infty|^p dx$$

which is not identically zero, and is contradiction. Hence, we have $t < 1$. By (4.2), we have

$$\begin{aligned}
m_\lambda &< \frac{1}{2} \int_{\mathbb{R}^3} (|\nabla \omega_\infty|^2 + V_\infty \omega_\infty^2) dx - \frac{\lambda}{4} \int_{\mathbb{R}^3} \rho_\infty \tilde{\Phi}[\omega_\infty] \omega_\infty^2 dx + \frac{1}{p} \int_{\mathbb{R}^3} Q_\infty |\omega_\infty|^p dx \\
&= I_\lambda^\infty(\omega_\infty) = m_\lambda^\infty.
\end{aligned}$$

Then there exists a positive radial ground state solution for the problem (1.1).

Acknowledgements

This research is supported by Shanghai Natural Science Foundation Project (No. 15ZR1429500), Shanghai Leading Academic Discipline Project (No. XTKX2012) and National Project Cultivate Foundation of USST (No. 13XGM05).

References

- [1] D'Aprile, T. and Mugnai, D. (2004) Solitary Waves for Nonlinear Klein-Gordon-Maxwell and Schrodinger-Maxwell Equations. *Proceedings of the Royal Society of Edinburgh: Section A*, **134**, 1-14.
- [2] D'Aprile, T. and Mugnai, D. (2004) Non-Existence Results for The coupled Klein-Gordon-Maxwell Equations. *Advanced Nonlinear Studies*, **4**, 307-322.
- [3] Ruiz, D. (2006) The Schrodinger-Poisson Equation under the Effect of a Nonlinear Local Term. *Journal of Functional Analysis*, **237**, 655-674. <http://dx.doi.org/10.1016/j.jfa.2006.04.005>
- [4] Ambrosetti, A. and Ruiz, D. (2008) Multiple Bound States for the Schrodinger-Poisson Problem. *Communications in*

- Contemporary Mathematics*, **10**, 391-404. <http://dx.doi.org/10.1142/S021919970800282X>
- [5] Ambrosetti, A. (2008) On Schrodinger-Poisson Problem Systems. *Milan Journal of Mathematics*, **76**, 257-274. <http://dx.doi.org/10.1007/s00032-008-0094-z>
- [6] Sanchez, O. and Soler, J. (2004) Long-Time Dynamics of the Schrodinger-Poisson-Slater System. *Journal of Statistical Physics*, **114**, 179-204. <http://dx.doi.org/10.1023/B:JOSS.0000003109.97208.53>
- [7] Mugnai, D. (2011) The Schrodinger-Poisson System with Positive Potential. *Communications in Partial Differential Equations*, **36**, 1099-1117. <http://dx.doi.org/10.1080/03605302.2011.558551>
- [8] Rabinowitz, P.H. (1992) On a Class of Nonlinear Schrodinger Equations. *Zeitschrift für angewandte Mathematik und Physik*, **43**, 270-291. <http://dx.doi.org/10.1007/BF00946631>
- [9] Cerami, G. and Vaira, G. (2010) Positive Solutions for Some Non-Autonomous Schrodinger-Poisson Systems. *Journal of Differential Equations*, **248**, 521-543. <http://dx.doi.org/10.1016/j.jde.2009.06.017>
- [10] Willem, M. (1996) *Minimax Theorems*, PNDEA Vol. 24. Birkhauser, Basel. <http://dx.doi.org/10.1007/978-1-4612-4146-1>
- [11] Talenti, G. (1976) Elliptic Equations and Rearrangements. *Annali della Scuola Normale Superiore di Pisa*, **3**, 697-718.

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

Other selected journals from SCIRP are listed as below. Submit your manuscript to us via either submit@scirp.org or [Online Submission Portal](#).

